

# Rational Function Multiplicative Coefficients

Michael Somos 11 Dec 2014

ms639@georgeown.edu

(draft version 11)

## 1 Rational generating functions of multiplicative sequences

Any numerical sequence has an associated generating function (GF). For example, the Fibonacci sequence is associated with GF  $x/(1-x-x^2)$ , a rational function of  $x$ . Consider a multiplicative sequence. That is,  $a(1) = 1$  and  $a(nm) = a(n)a(m)$  for all positive integers  $n$  and  $m$  relatively prime to each other. Can its GF  $f(x) = a(1)x + a(2)x^2 + a(3)x^3 + \dots$  ever be rational? The answer is yes if  $f(x) = x/(1-x)$  and  $a(n) = 1$  if  $n > 0$ . This is the simplest example where  $a(n)$  is non-zero for all  $n > 0$ . Another is  $f(x) = x/(1-x^2)$  and  $a(n) = 1$  if  $n > 0$  is odd and  $a(n) = 0$  otherwise. Now consider the rational function and its power series expansion

$$f(x) = x(1-x)^{e_1}(1-x^2)^{e_2} = x - e_1x^2 + ((e_1^2 - e_1)/2 - e_2)x^3 + \dots$$

where  $e_1$  and  $e_2$  are integers. A search finds that  $f(x)$  is the GF of a multiplicative sequence for 11 pairs of integers  $[e_1, e_2]$  as follows:

$$[-4, 1], [-2, 0], [-1, 0], [-1, 1], [0, -1], [0, 0], [0, 1], [1, -1], [1, 0], [2, -2], [4, -3].$$

The multiplicative integer sequences for these pairs are of a simple form. Some algebra is enough to prove that this list is complete. Allowing more factors in  $f(x)$  increases the difficulty of search and algebraic proof.

## 2 Conjecture 1

Conjecture 1: there is a finite set of rational functions of the form

$$f(x) = x(1-x)^{e_1}(1-x^2)^{e_2}\dots(1-x^n)^{e_n}$$

for some integers  $e_1, \dots, e_n$  which are the GF for multiplicative integer sequences provided we exclude some infinite families which are predictable. One example infinite family is

$$f(x) = x(1-x^{n-1}) = x - x^n$$

where  $n > 1$ . Also, if and only if  $n = p^k, n > 1$  and  $p$  is prime then

$$f(x) = x(1-x)^{-1}(1-x^{n-1})(1-x^n)^{-1}$$

is multiplicative. Note that  $f(x)$  is in the set when  $-f(-x)$  is since  $(1+x) = (1-x^2)/(1-x)$  and so on.

### 3 Homogeneous generalization of multiplicative sequences

Now assume  $a(0)$  and  $a(1)$  are nonzero. For example, consider the sequence  $a(n)$  with its GF

$$g(x) = (1-x)^2/(1-x^2) = 1 - 2x + 2x^2 - 2x^3 + \dots$$

Then  $a(1)a(nm) = a(n)a(m)$  for all positive integers  $n$  and  $m$  relatively prime to each other. This is a homogeneous generalization of multiplicative sequences. As in the first section, but without a factor of  $x$ , consider

$$g(x) = (1-x)^{e_1}(1-x^2)^{e_2} = 1 - e_1x + ((e_1^2 - e_1)/2 - e_2)x^2 + \dots$$

where  $e_1$  and  $e_2$  are integers. A search finds that  $g(x)$  is the GF of a homogeneous multiplicative sequence for 10 pairs of integers  $[e_1, e_2]$  as follows:

$$[-4, 2], [-2, 1], [-2, 2], [-1, 0], [-1, 1], [1, -1], [1, 0], [2, -1], [2, 0], [4, -2].$$

Again, algebra is enough to prove the list is complete.

### 4 Conjecture 2

Conjecture 2: there is a finite set of rational functions of the form

$$g(x) = (1-x)^{e_1}(1-x^2)^{e_2} \dots (1-x^n)^{e_n}$$

for some integers  $e_1, \dots, e_n$  which are the GF for homogeneous multiplicative integer sequences provided we exclude some infinite families which are predictable. For example,

$$g(x) = (1-x)^{-1}(1-x^n)^{-1}(1-x^{n+1})$$

is homogeneous multiplicative if and only if  $n = p^k$ ,  $n > 1$  and  $p$  is prime. Note that  $g(x)$  is in the set when  $g(-x)$  is.

### 5 Further Work

The rational functions in the two conjectures have applications related to Ramanujan's Lambert series. A study of rational functions with poles only at roots of unity appeared in 2003 by Juan B. Gil and Sinai Robins who defined a Hecke operator on power series. Kyoji Saito studied cyclotomic functions related to eta-products in 2001. Rational functions of a simple form having multiplicative coefficients is related to a paper on Multiplicative  $\eta$ -Quotients by Yves Martin in 1996.